

ASTHMA ACTION PLAN

Take me when you visit your doctor



photo (optional)

Patient name:

Plan date: Review date:

Doctor details:

EMERGENCY CONTACT

Name:

Phone:

Relationship:

WELL CONTROLLED is all of these...

- needing reliever medication no more than 2 days/week
- no asthma at night
- no asthma when I wake up
- can do all my activities

Peak Flow reading (if used) above

TAKE preventer

name

day / night puffs/inhalations

Use my preventer, even when well controlled Use my spacer with my puffer

TAKE reliever

name

puffs/inhalations as needed puffs/inhalations 15 minutes before exercise

Always carry my reliever medication

FLARE-UP is any of these...

- needing reliever medication more than usual OR days/week
- woke up overnight with asthma
- had asthma when I woke up
- can't do all my activities

Peak Flow reading (if used) between and

my triggers and symptoms

TAKE preventer

name

day / night puffs/inhalations for days then back to **well controlled** dose

TAKE reliever

name puffs/inhalations as needed

START other medication

name dose for days

MAKE an appointment to see my doctor this week

SEVERE is any of these...

- reliever medication not lasting 3 hours
- woke up frequently overnight with asthma
- had asthma when I woke up
- difficulty breathing

Peak Flow reading (if used) between and

my triggers and symptoms

TAKE preventer

name

day / night puffs/inhalations for days then back to **well controlled** dose

TAKE reliever

name puffs/inhalations as needed

START other medication

name dose for days

MAKE an appointment to see my doctor TODAY

If unable to see my doctor, visit a hospital

If unable to see my doctor/hospital:

START other medication

name dose for days

EMERGENCY is any of these...

- reliever medication not working
- can't speak a full sentence
- extreme difficulty breathing
- feel asthma is out of control
- lips turning blue

Peak Flow reading (if used) below

1 **CALL AMBULANCE NOW**
Dial Triple Zero (000)

2 **START ASTHMA FIRST AID**
Turn page for Asthma First Aid

ASTHMA FIRST AID

Blue/Grey Reliever

Airomir, Asmol, Ventolin or Zempreon and Bricanyl

Blue/grey reliever medication is unlikely to harm, even if the person does not have asthma



DIAL TRIPLE ZERO (000) FOR AN AMBULANCE IMMEDIATELY IF THE PERSON:

- is not breathing
- suddenly becomes worse or is not improving
- is having an asthma attack and a reliever is not available
- is unsure if it is asthma
- has a known allergy to food, insects or medication and has **SUDDEN BREATHING DIFFICULTY, GIVE ADRENALINE AUTOINJECTOR FIRST (if available), even if there are no skin changes, then use a reliever**

1



SIT THE PERSON UPRIGHT

- Be calm and reassuring
- Do not leave them alone

2



GIVE 4 SEPARATE PUFFS OF RELIEVER PUFFER

- Shake puffer
- Put 1 puff into spacer
- Take 4 breaths from spacer
 - Repeat until 4 puffs have been taken



If using **Bricanyl**, give 2 separate inhalations (5 years or older)

If you don't have a spacer handy in an emergency, take 1 puff as you take 1 slow, deep breath and hold breath for as long as comfortable. **Repeat** until all puffs are given

3



WAIT 4 MINUTES

- If breathing does not return to normal, give 4 more separate puffs of reliever as above



Bricanyl: Give 1 more inhalation

IF BREATHING DOES NOT RETURN TO NORMAL

4



DIAL TRIPLE ZERO (000)

- Say 'ambulance' and that someone is having an asthma attack
- Keep giving 4 separate puffs every 4 minutes until emergency assistance arrives



Bricanyl: Give 1 more inhalation every 4 minutes until emergency assistance arrives